

A-31

Activity and Ordering of Mixed Phosphatidylethanolamine/Dihydrocholesterol Monolayers

Kathleen D. Cao¹, Luka Pocivavsek¹, Niels Holten-Andersen¹, Stephanie Harmon², Jaroslaw Majewski³, Mati Meron⁴, Binhua Lin⁴, and Ka Yee C. Lee¹

¹Department of Chemistry, Institute for Biophysical Dynamics, and The James Franck Institute, University of Chicago, Chicago, IL 60637

²Biological, Chemical, and Physical Sciences Department, Illinois Institute of Technology, Chicago, IL 60616

³Manuel Lujan Neutron Scattering Center, Los Alamos National Laboratory, Los Alamos, NM 87545

⁴The James Franck Institute and Center for Advanced Radiation Sources, University of Chicago, Chicago, IL 60637

Cholesterol is thought to be important for the structure and assembly of lipid rafts, and its interaction with other membrane lipids has been a topic of great research interest. The interactions between 1,2-dimyristoyl-sn-glycero-3-phosphoethanolamine (DMPE) and dihydrocholesterol (Dchol) in Langmuir monolayers are probed using fluorescence microscopy (FM), beta-cyclodextrin (CD) desorption assays, and grazing-incidence x-ray diffraction (GIXD). Similar to our previous results for 1,2-dimyristoyl-sn-glycero-3-phosphocholine (DMPC)/Dchol monolayers [1], FM and CD assays show two regimes for the DMPE/Dchol system. Short-ranged lateral ordering was observed using GIXD that was also consistent with our recent work on sphingomyelin (SM)/Dchol monolayers [2]. We investigate how the smaller headgroup of DMPE effects Dchol chemical activity and lateral structure compared to monolayers of Dchol with DMPC or SM.

1. *Biophys. J.*, **93**, 2038–2047, (2007).

2. *Phys. Rev. Lett.*, **103**, 028103, (2009).